

THE LANCET Oncology

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed.
We post it as supplied by the authors.

Supplement to: Pinato DJ, Tabernero J, Bower M, et al. Prevalence and impact of COVID-19 sequelae on treatment and survival of patients with cancer who recovered from SARS-CoV-2 infection: evidence from the OnCovid retrospective, multicentre registry study. *Lancet Oncol* 2021; published online Nov 3. [http://dx.doi.org/10.1016/S1470-2045\(21\)00573-8](http://dx.doi.org/10.1016/S1470-2045(21)00573-8).


Prevalence and impact of COVID-19 sequelae on treatment and survival of patients with cancer who recovered from SARS-Cov-2 infection: evidence from the OnCovid retrospective, multicentre registry study

APPENDIX


CONTENTS

Supplementary Figure 1	p. 2
Post COVID-19 survival estimate.	
Supplementary Figure 2	p. 3
COVID-19 Sequelae	
Supplementary Figure 3	p. 4
Post COVID-19 survival estimates.	
Supplementary Figure 4	p. 5
Post COVID-19 survival estimates.	
Supplementary Figure 5	pp. 6-7
SACT resumption pathways.	
Supplementary Table 1	p. 8
Detailed comorbidities.	
Supplementary Methods	pp. 8-9
Supplementary Table 2	p. 10
Patient disposition across participating centres.	
Supplementary Table 3	p. 11
Patients and disease characteristics	
Supplementary Table 4	p. 12
Multivariable analysis for post COVID-19 survival (SACT pathway)	
Supplementary Table 5	p. 13
Patients and disease characteristics according to the SARS-CoV-2 antibody status	
OnCovid Study Group: Investigators	p. 14
Figure 2 legend	p. 17


Supplementary Figure 1. Kaplan Meier survival estimate of post COVID-19 survival (days). The median post COVID-19 survival (929 patients) was not reached with 143 events




Supplementary Figure 2. COVID-19 sequelae. Overall 234 patients experienced at least 1 COVID-19 sequelae. Respiratory sequelae: 116 patients (49.6%), including chronic cough in 15 patients (6.4%), residual dyspnoea/shortness of breath in 79 patients (33.8%) and other respiratory sequelae in 25 patients (10.7%); chronic fatigue: 96 patients (41.0%); weight loss: 13 patients (5.5%), neuro-cognitive disfunctions (including dysgeusia/dysosmia): 17 patients (7.3%); non-respiratory organ disfunction: 4 patients (1.7%); others: 43 patients (18.4%).



Supplementary Figure 3. A) Kaplan Meier survival estimate of post COVID-19 survival (days) according to the time from cancer diagnosis to post COVID-19 reassessment (not reached, 825 patients included (HR 1.03, 95%CI: 0.72-1.47; p=0.8570). **B)** Kaplan Meier survival estimate of post COVID-19 survival (days) according to the time from COVID-19 diagnosis to post COVID-19 reassessment. The median post COVID-19 survival (929 patients) was not reached with 143 events HR 1.13, 95%CI: 0.81-1.22; p=0.4431) (929 patients included)



Supplementary Figure 5. A) Detailed underlying causes for SACT permanent cessation (70 patients) and regimen adjustments (178 patients). Reason to permanent discontinuation (31 patients): 19 (61.3%) performance status (PS) deterioration, 9 (29%) disease progression during COVID-19, 3 (9.7%) residual organ disfunction. Dose/regimen adjustments: 89 (50%) avoid potential immune suppression, 46 (25.8%) avoid hospital attendance, 14 (7.9%), avoid intravenous administration, 34 (19.1%) avoid other adverse events, 36 (20.2%) other reasons. **B)** Histogram reporting the paired prevalence of COVID-19 sequelae according to post COVID-19 SACT resumption pathways (the Pearson χ^2 test was used). COVID-19 sequelae (yes vs no): SACT unchanged: 21/218 (9.6%), dose/regimen adjusted: 26/178 (14.6%), permanent cessation: 16/70 (22.9%).




Supplementary Figure 5C. Histogram with paired prevalence of SACT permanent cessation and dose/regimen adjustments according to the primary tumour category.

Permanent cessation plus dose/regimen adjustments: breast 52/122 (42.6%), gastro-intestinal 52/83 (62.7%), gynaecological/genito-urinary 31/81 (38.3%), haematological 46/68 (67.6%), thoracic 37/62 (59.7%), others 29/47 (61.7%). Chi-squared: 23.8, DF: 5; p=0.0002.

Permanent cessation (dose/regimen adjustments excluded): breast 10/80 (12.5%), gastro-intestinal 23/54 (42.6%), gynaecological/genito-urinary 8/58 (13.8%), haematological 6/28 (21.4%), thoracic 15/40 (37.5%), others 8/26 (30.8%). Chi-squared: 23.7, DF: 5; p=0.0002.

Dose regimen adjustments (permanent cessation excluded): breast 42/112 (37.5%), gastro-intestinal 29/60 (48.3%), gynaecological/genito-urinary 23/73 (31.5%), haematological 40/62 (64.5%), thoracic 22/47 (46.8%), others 21/39 (53.8%). Chi-squared: 19.0, DF: 5; p=0.0019.



Supplementary Table 1: Detailed comorbidities.

Comorbidities	
Ischaemic heart disease	79 (7.8)
Dementia	33 (3.3)
Diabetes	183 (18.0)
Immunosuppression	23 (2.3)
Congestive cardiac failure	14 (1.4)
Liver impairment	14 (1.4)
Hypertension	420 (41.4)
Peripheral vascular disease	29 (2.9)
Cardiovascular disease	110 (10.8)
Cerebro-vascular disease	44 (4.3)
Chronic pulmonary disease	120 (12.99)
Chronic kidney disease	58 (5.7)
Obesity (BMI > 30)	29 (2.9)
Others	249 (24.5)

Supplementary Methods

OnCovid (NCT04393974) is an active European registry study that, since the beginning of the pandemic, has collected consecutive patients fulfilling the following inclusion criteria: 1) age ≥18 years; 2) diagnosis of SARS-CoV-2 infection confirmed by RT-PCR of a nasopharyngeal swab; 3) history of solid or hematologic malignancy, at any time during the patients' past medical history, either active or in remission at the time of COVID-19 diagnosis. Patients with a history of non-invasive/premalignant lesions or with low malignant potential (i.e., basal cell carcinoma of the skin, non-invasive carcinoma in situ of the cervix, ductal carcinoma in situ) were excluded. For hematologic malignancies, only patients with a history of oncologic diseases with defined malignant behavior (lymphoma, leukaemia, multiple myeloma) were included.

OnCovid was granted central approval by the United Kingdom Health Research Authority (20/HRA/1608) and by the corresponding research ethics committees at each participating institution. Core study data were collated from electronic medical records into a case report form designed using the Research Electronic Data Capture software (REDCap, Vanderbilt University, Nashville, TN, USA). Multi-site access and data curation was coordinated by the Medical Statistics Unit in Novara, Italy.

This OnCovid registry focused on post COVID-19 outcomes among COVID-19 survivors who underwent a formal clinical reassessment at the participating institutions.

COVID-19 sequelae were defined as any residual symptoms and/or measurable organ disfunction attributable to COVID-19 according to investigators. Patterns of SACT resumption were assessed among patients who were on systemic anti-cancer therapy (SACT) within 4 weeks of COVID-19 diagnosis.

Accounting for the unbalanced distribution of patient/disease-related features across the subgroups we adopted a fixed multivariable regression model, adjusting all survival estimates for clinical characteristics already known to influence clinical outcomes in patients with COVID-19 and cancer. Included covariates were incorporated following a clinical prioritization. In addition, all of them revealed to be associated with both the short-term and medium-term outcomes at the latest updated of the registry including 1392 patients (Pinato DJ, et al. Eur J of Cancer 2021).

The following key variables of interest were used as covariates within the survival analysis:

- Gender (male vs female),
- Age (≥65 vs < 65 years),
- Number of co-morbidities (0-1 vs ≥ 2),
- Primary tumour (clustered as: breast, gastro-intestinal, gynaecological/genito-urinary, haematological, thoracic, and others),
- Tumour stage (defined as advanced vs non-advanced). In details, we defined as “advanced” stage any patient with distant metastatic disease, to differentiate them from “non-advanced” patients. Disease-specific criteria (i.e. Rai, Binet criteria etc.) were utilised as appropriate to define advanced haematological malignancies.
- Tumour status (presence of active vs non-active disease), on the basis of disease-specific criteria (radiologic, clinical, biochemical/haematological depending on disease type).

- Receipt of any anticancer therapy within 4 weeks of SARS-CoV-2 infection, including non-systemic therapy (yes vs no),
- Experience of at least one COVID-19 complications including acute respiratory failure, ARDS, kidney injury, secondary infections, sepsis, septic shock, acute cardiac injury, acute liver injury and others (yes vs no);
- Receipt of any COVID-19 specific therapy, including antivirals, antimalarials, antibiotics, corticosteroids and others (yes vs no),
- Hospitalization requirement (pre-existent/due to COVID-19 vs not required).

In order to provide an estimate of COVID-19 sequelae distribution across different age-groups and to account for their possible relationship with COVID-19 severity, we described sequelae proportion, rate of complicated COVID-19 and hospital admission due to COVID-19, according to 6 age-categories: <40 years old, 40-49 years old, 50-59 years old, 60-69 years old, 70-79 years old, \geq 80 years old.

Being an observational study, decisions on initiating, resuming or discontinuing therapy were dictated by the discretion of the treating clinicians. The overarching subgrouping of tumour related features according to stage, activity and therapy has been consistently utilised in all the publications from our registry (Cancer Discov. 2020 Jul 31;10(10):1465–74, Eur J Cancer. 2021 Jun;150:190-202, J Immunother Cancer. 2021 Mar;9(3):e002277, Cancers. 2020 Jul 8;12(7):1841) and was made necessary by the wide heterogeneity oncological diagnoses included in the registry.

Additional variables reported in the descriptive analysis were: smoking history (former/current smokers vs never smokers), systemic anticancer therapy (chemotherapy, immune checkpoint inhibitors, other targeted agents and endocrine therapy), median time (in days) from symptoms to SARS-CoV-2 infection diagnosis, COVID-19 therapy details (antibiotics, antimalarials, antivirals, IL-6 inhibitors, corticosteroids and others), intensive care admission, mechanical ventilation requirements, and oxygen therapy.

Oncological and disease specific variables were collected at baseline, defined at the moment of diagnosis of SARS-Cov-2 by PCR test. Characteristics of severity, complications and therapy against COVID-19 were collected throughout the observation period until full clinical resolution of COVID-19 or patients' mortality.

Patient observation time started from date of first PCR/SARS-CoV-2 infection confirmation until patient death or loss to follow-up. Being a retrospective, observational study, the entirety of the OnCovid cohort was followed up at intervals dictated by the routine clinical practice in each participating institutions, as deemed clinically indicated by the treating physicians. All-cause of mortality was retrieved and validated by investigators at each centre by accessing patients' electronic medical records and death certificates.

Patients were lost to follow-up when for any reason failed to attend planned follow-up appointments scheduled by the treating clinicians. Given the pragmatic nature of this registry, based on standard of care clinical practice, we could not accurately reconstruct the reasons to explain why a proportion of patients did not attend for follow-up. To avoid incurring into bias, by mislabelling patients that were lost to follow-up as potentially deceased, we decided to exclude all patients with incomplete/missing follow up data to preserve the integrity of our results.

We also performed an ancillary analysis evaluating post COVID-19 anti-SARS-CoV-2-S (Spike) IgG seroprevalence. To define patient/disease characteristics possibly related to seroconversion, their distribution according to the antibody status were described. Serum SARS-CoV-2 antibodies were tested in clinical practice at participating institution and seroconversion has been defined as the presence of positive anti-SARS-CoV-2-S (Spike) IgG antibody test after COVID-19 resolution.

Supplementary Table 2: Patient disposition across participating centres. PI: principal investigator.

Institution	Patients		Site PI
	N	%	
Chelsea and Westminster Hospital, London (UK)	361	13.7%	Mark Bower
Vall d'Hebron University Hospital, Barcelona (Spain)	294	11.2%	Josep Tabernero
University College London, London (UK)	209	7.9%	Diego Ottaviani
Ospedale Maggiore della Carità, Novara (Italy)	207	7.9%	Alessandra Gennari
Institut Gustave Roussy, Villejuif (France)	187	7.1%	Emeline Colomba
Guy's and St Thomas' NHS Foundation Trust, London (UK)	150	5.7%	Saoirse Dolly
Barts Health NHS Trust, London (UK)	145	5.5%	Nikolaos Diamantis
Humanitas Cancer Centre, Milan (Italy)	139	5.3%	Alexia Bertuzzi
ICO L'Hospitalet, L'Hospitalet de Llobregat, Barcelona (Spain)	123	4.7%	Ramon Salazar
Ospedale Papa Giovanni XXIII, Bergamo (Italy)	107	4.1%	Alberto Zambelli
Policlinico San Matteo, Pavia (Italy)	67	2.5%	Gianpiero Rizzo
ICO Badalona (Spain)	61	2.3%	Eudald Felip
IRCCS AOU San Martino, Genova (Italy)	56	2.2%	Matteo Lambertini
ICO Girona (Spain)	56	2.1%	Joan Brunet
Hospital Clinic, Barcelona (Spain)	53	2.0%	Aleix Prat
Ospedale Antonio e Biagio e Cesare Arrigo, Alessandria (Italy)	53	2.0%	Antonio Maconi
Imperial College London, London (UK)	46	1.7%	David J Pinato
Velindre Cancer Centre, Cardiff (UK)	34	1.3%	John Chester
Manresa (Spain)	33	1.3%	Clara Martinez-Vila
Institut Jules Bordet, Brussels (Belgium)	30	1.1%	Angela Loizidou
University of L'Aquila, L'Aquila (Italy)	30	1.1%	Corrado Ficorella
Azienda Ospedaliera Spedali Civili, Brescia (Italy)	26	1.0%	Salvatore Grisanti
Azienda Istituti Ospitalieri di Cremona, Cremona (Italy)	25	0.9%	Daniele Generali
Università Campus Bio-Medico, Rome (Italy)	20	0.8%	Buno Vincenzi
University of Munich (Germany)	19	0.7%	Nadia Harbeck
Hospital Universitario 12 de Octubre, Madrid (Spain)	15	0.6%	Ana Sanchez de Torre
Azienda Ospedaliera S Maria, Terni (Italy)	14	0.5%	Annalisa Guida
Ospedali Riuniti di Ancona, Università Politecnica delle Marche (Italy)	14	0.5%	Rossana Berardi
Istituto Europeo di Oncologia, Milano (Italy)	10	0.4%	Paola Queirolo
Fondazione Poliambulanza Istituto Ospedaliero, Brescia (Italy)	10	0.4%	Michela Libertini
Istituto Tumori, Milan (Italy)	8	0.3%	Rossella Bertulli
Careggi University Hospital, Florence (Italy)	8	0.3%	Francesca Mazzoni
Santa Maria Goretti Hospital, Latina (Italy)	8	0.3%	Federica Zoratto
Azienda Ospedaliera S. Andrea, Rome (Italy)	8	0.3%	Raffaele Giusti
Palma de Mallorca Hospital, Palma de Mallorca, (Spain)	4	0.2%	Maria Iglesias
University of Bari 'Aldo Moro', Bari (Italy)	4	0.2%	Marco Tucci
Total	2634	100.0%	

Supplementary Table 3: Distribution of baseline patients, tumour and COVID-19 characteristics among the overall entered population and among patients who underwent a formal clinical re-assessment at participating institutions.

Characteristic	Overall Population	Post-COVID-19 analysis
	N = 2634 (%)	N = 1557 (%)
Sex		
Male	1390 (52.8)	751 (48.3)
Females	1240 (47.2)	803 (51.7)
Missing	4	3
Age		
<65 years	1083 (41.3)	788 (50.8)
≥65 years	1538 (58.7)	763 (49.2)
Missing	13	6
Comorbidities		
0-1	1414 (53.7)	962 (61.8)
≥2	1220 (46.3)	595 (38.2)
Smoking history		
Never smokers	1128 (51.9)	732 (55.7)
Former/current smokers	1044 (48.1)	583 (44.3)
Missing	462	242
Primary Tumour		
Breast	493 (18.9)	360 (23.4)
Gastrointestinal	476 (18.2)	255 (16.5)
Gynaecological/Genito-Urinary	530 (20.3)	297 (19.3)
Haematological	357 (13.7)	218 (14.1)
Thoracic	375 (14.4)	188 (12.2)
Others	382 (14.6)	223 (14.5)
Missing	21	16
Tumour stage		
Local/loco-regional	1237 (53.96)	741 (50.7)
Advanced	1244 (46.04)	720 (49.3)
Missing	153	96
Tumour status at COVID-19 diagnosis		
Remission/non measurable disease	860 (33.3)	520 (33.9)
Active malignancy	1723 (66.7)	1015 (66.1)
Missing	51	22
Anticancer therapy at COVID-19 diagnosis		
No	1213 (48.2)	819 (54.5)
Yes	1305 (51.8)	685 (45.5)
Missing	116	53
COVID-19 therapy		
No	1129 (42.9)	707 (45.4)
Yes	1505 (57.1)	850 (54.6)
Antibiotics	1242 (47.1)	650 (41.7)
Antimalarials	615 (23.3)	394 (25.3)
Antiviral	400 (15.2)	231 (14.8)
IL-6 Inhibitors	93 (3.5)	59 (3.8)
Corticosteroids	489 (18.6)	248 (15.9)
Other	162 (6.2)	104 (6.7)
Complicated COVID-19		
No	1554 (59.0)	1154 (74.1)
Yes	1080 (41.0)	403 (25.9)
Hospitalization		
Not required	593 (22.7)	519 (33.5)
Required	1387 (53.1)	711 (45.9)
Pre-existing	633 (24.2)	318 (20.5)
Missing	21	9
Intensive care admission		
No	1649 (84.5)	880 (87.7)
Yes	273 (14.0)	117 (11.7)
Indicated but not admitted	30 (1.5)	6 (0.6)
Missing	682	554
Mechanical ventilation		
No	2117 (88.6)	1343 (93.9)
Yes	273 (11.4)	87 (6.1)
Missing	244	127
Oxygen therapy		
No	1208 (48.7)	930 (62.8)
Yes	1272 (51.3)	551 (37.2)
Missing	154	76

Supplementary Table 4: Fixed multivariable regression model for post COVID-19 survival including patients on SACT at COVID-19 diagnosis (309 patients included). HR: hazard ratio; CI: confidence interval.

Oncology therapeutic pathway	Multivariable analysis
	HR 95%CI; p-value
SACT	
Unchanged	1
Dose/regimen adjusted	0.84 (0.35-2.02)
Permanent cessation	3.53 (1.45-8.59)
Sex	
Female	1
Male	0.72 (0.34-1.50)
Age	
<65 years	1
≥65 years	1.27 (0.64-2.52)
Comorbidities	
0-1	1
≥2	1.46 (0.71-3.03)
Primary tumour	
Breast	2.09 (0.31-14.07)
Gastrointestinal	3.94 (0.80-19.30)
Gynaecological/Genito-Urinary	3.79 (0.72-19.86)
Thoracic	3.27 (0.60-17.79)
Other	6.15 (1.19-31.65)
Haematological	1
Tumour stage	
Local/loco-regional	1
Advanced	2.49 (0.90-6.86)
Tumour status	
Remission/non measurable disease	1
Active malignancy	7.12 (0.90-56.33)
COVID-19 complications	
0	1
≥1	2.92 (1.25-6.78)
COVID-19 therapy	
No	1
Yes	0.71 (0.35-1.45)
Hospitalization	
Not- Required	1
Required due to COVID-19	1.27 (0.40-4.06)
Pre-existing	3.27 (1.03-10.38)
COVID-19 sequelae	
No	1
Yes	1.21 (0.51-2.88)

Supplementary Table 5: Distribution of baseline patients, tumour and COVID-19 characteristics according to the post COVID-19 SARS-CoV-2 antibody status. *Within 4 weeks of COVID-19 diagnosis.

	Negative SARS-CoV-2 antibodies	Positive SARS-CoV-2 antibodies	P value
	N = 32 (%)	N = 318 (%)	
Sex			
Male	10 (31.2)	141 (44.3)	0.15
Females	22 (68.7)	177 (55.7)	
Age			
<65 years	19 (59.4)	176 (55.5)	0.67
≥65 years	13 (40.6)	141 (44.5)	
Comorbidities			
0-1	24 (75.0)	210 (66.0)	0.31
≥2	8 (25)	108 (34.0)	
Smoking history			
Never smokers	20 (71.4)	163 (58.6)	0.18
Former/current smokers	8 (28.6)	115 (41.4)	
Missing	4	40	
Primary Tumour			
Breast	9 (28.1)	106 (33.7)	0.35
Gastrointestinal	3 (9.4)	42 (13.3)	
Gynaecological/Genito-Urinary	8 (25.0)	44 (14.0)	
Haematological	8 (25.0)	57 (18.1)	
Thoracic	3 (9.4)	31 (9.8)	
Others	1 (3.1)	35 (11.1)	
Missing	0	3	
Tumour stage			
Local/loco-regional	14 (46.7)	138 (46.8)	0.99
Advanced	16 (53.3)	157 (53.2)	
Missing	2	23	
Tumour status at COVID-19 diagnosis			
Remission/non measurable disease	9 (29.0)	116 (37.2)	0.36
Active malignancy	22 (71.0)	196 (62.8)	
Missing	1	6	
Anticancer therapy at COVID-19 diagnosis*			
No	22 (68.7)	191 (62.4)	0.48
Yes	10 (31.2)	115 (37.6)	
Missing	0	12	
COVID-19 therapy			
No	19 (59.4)	119 (37.4)	0.02
Yes	13 (40.6)	199 (62.6)	
Antibiotics	9 (28.1)	140 (44.0)	0.08
Antimalarials	9 (28.1)	115 (36.2)	0.36
Antiviral	4 (12.5)	84 (26.4)	0.08
IL-6 Inhibitors	2 (6.2)	19 (6.0)	0.95
Corticosteroids	5 (15.6)	62 (19.5)	0.59
Other	-	4 (1.3)	0.52
Complicated COVID-19			
0	26 (81.2)	238 (74.8)	0.42
≥ 1	6 (18.8)	80 (25.2)	
COVID-19 sequelae			
No	28 (87.5)	267 (84.0)	0.60
Yes	4 (12.5)	51 (16.0)	
Hospitalization			
Not required	15 (46.9)	100 (31.7)	0.21
Required	14 (43.7)	169 (53.7)	
Pre-existing	3 (9.4)	46 (14.6)	
Missing	0	3	

OnCovid Study group: Investigators

NAME	SURNAME	INSTITUTION
David J	Pinato	Imperial College London, London, UK
Joanne S	Evans	Imperial College London, London, UK
Judith	Swallow	Imperial College London, London, UK
Alessio	Cortellini	Imperial College London, London, UK
Chris	Chung	Imperial College London, London, UK
Meera	Patel	Imperial College London, London, UK
Gino	Dettorre	Imperial College London, London, UK
Diego	Ottaviani	University College London, London UK
Amani	Chowdhury	University College London, London UK
Eve	Merry	University College London, London UK
Neha	Chopra	University College London, London UK
Alvin JX	Lee	University College London, London UK
Christopher CT	Sng	University College London, London UK
Yien Ning Sophia	Wong	University College London, London UK
Myria	Galazi	University College London, London UK
Sarah	Benafif	University College London, London UK
Palma	Dileo	University College London, London UK
Irina	Earnshaw	University College London, London UK
Grisma	Patel	University College London, London UK
Anjui	Wu	University College London, London UK
Alasdair	Sinclair	University College London, London UK
Gehan	Soosaipillai	University College London, London UK
Mark	Bower	Chelsea and Westminster Hospital, London UK
Alessia	Dalla Pria	Chelsea and Westminster Hospital, London UK
Rachel	Sharkey	Chelsea and Westminster Hospital, London UK
Thomas	Newsom-Davis	Chelsea and Westminster Hospital, London UK
Dolly	Saorise	Guy's and St Thomas' NHS Foundation Trust, London UK
Ailsa	Sita-Lumsde	Guy's and St Thomas' NHS Foundation Trust, London UK
Krishnie	Srikandarajah	Guy's and St Thomas' NHS Foundation Trust, London UK
Eleanor	Jones	Guy's and St Thomas' NHS Foundation Trust, London UK
Mieke	Van Hemelrijck	King's College London, London UK
Charlotte	Moss	King's College London, London UK
Beth	Russell	King's College London, London UK
Nikolaos	Diamantis	Barts Health NHS Trust, London UK
Uma	Mukherjee	Barts Health NHS Trust, London UK
Amanda	Jackson	Velindre Cancer Centre, Cardiff, UK
John	Chester	Velindre Cancer Centre, Cardiff, UK
Angela	Loizidou	Institut Jules Bordet, Brussels, Belgium
Martine	Piccart	Institut Jules Bordet, Brussels, Belgium
Fanny	Pommeret	Institut Gustave Roussy, Villejuif, France
Johann	Colomba	Institut Gustave Roussy, Villejuif, France
Emeline	Colomba-Blameble	Institut Gustave Roussy, Villejuif, France
Aleix	Prat	Hospital Clinic Barcelona, Barcelona, Spain
Claudia A	Cruz	Hospital Clinic Barcelona, Barcelona, Spain
Roxana	Reyes	Hospital Clinic Barcelona, Barcelona, Spain
Elia	Segui	Hospital Clinic Barcelona, Barcelona, Spain

Javier	Marco-Hernández	Hospital Clinic Barcelona, Barcelona, Spain
Josep	Tabernero	Vall dHebron University Hospital, Barcelona, Spain
Juan	Aguilar-Company	Vall dHebron University Hospital, Barcelona, Spain
Isabel	Ruiz-Camps	Vall dHebron University Hospital, Barcelona, Spain
Laura	Fox	Vall dHebron University Hospital, Barcelona, Spain
David	Garcia Illescas	Vall dHebron University Hospital, Barcelona, Spain
Nadia	Saoudi	Vall dHebron University Hospital, Barcelona, Spain
Oriol	Mirallas	Vall dHebron University Hospital, Barcelona, Spain
Elisa	Roldán	Vall dHebron University Hospital, Barcelona, Spain
Joan	Brunet	Catalan, Institute of Oncology, Girona, Spain
Anna	Carbó	Catalan, Institute of Oncology, Girona, Spain
Carmen	Carmona	Catalan, Institute of Oncology, Girona, Spain
Isabel	Garcia	Catalan, Institute of Oncology, Girona, Spain
Raquel	Liñan	Catalan, Institute of Oncology, Girona, Spain
Ariadna	Roqué Lloveras	Catalan, Institute of Oncology, Girona, Spain
Nadia	Harbeck	University Hospital of Munich, Germany
Rachel	Wuerstlein	University Hospital of Munich, Germany
Ricard	Mesia	Institut Català dOncologia Badalona, Spain
Eudald	Felip	Institut Català dOncologia Badalona, Spain
Ramon	Salazar	ICO LHospitalet, LHospitalet de Llobregat, Barcelona, Spain
Anna	Sureda	ICO LHospitalet, LHospitalet de Llobregat, Barcelona, Spain
Clara	Maluquer	ICO LHospitalet, LHospitalet de Llobregat, Barcelona, Spain
Alessandra	Gennari	University of Piemonte Orientale, Novara, Italy
Federica	Biello	University of Piemonte Orientale, Novara, Italy
Francesca	D'Avanzo	University of Piemonte Orientale, Novara, Italy
Gianluca	Gaidano	University of Piemonte Orientale, Novara, Italy
Riccardo	Bruna	University of Piemonte Orientale, Novara, Italy
Andrea	Patriarca	University of Piemonte Orientale, Novara, Italy
Daniela	Ferrante	University of Piemonte Orientale, Novara, Italy
Lorenza	Scotti	University of Piemonte Orientale, Novara, Italy
Marco	Krengly	University of Piemonte Orientale, Novara, Italy
Paolo	Pedrazzoli	University of Pavia, Pavia, Italy
Gianpiero	Rizzo	Fondazione IRCCS Policlinico San Matteo, Pavia, Italy
Alexia	Bertuzzi	Humanitas Cancer Center, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy
Sabrina	Rossi	Humanitas Cancer Center, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy
Andrea	Marrari	Humanitas Cancer Center, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy

Armando	Santoro	Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy; Humanitas Cancer Center, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy
Lorenza	Rimassa	Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy; Humanitas Cancer Center, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy
Antonio	Maconi	Ospedale Antonio e Biagio e Cesare Arrigo, Alessandria, Italy
Marta	Betti	Ospedale Antonio e Biagio e Cesare Arrigo, Alessandria, Italy
Bruno	Vincenzi	Università Campus Bio-Medico, Roma, Italy
Giuseppe	Tonini	Università Campus Bio-Medico, Roma, Italy
Alberto	Zambelli	ASST Papa Giovanni XXIII, Bergamo, Italy
Carlo	Tondini	ASST Papa Giovanni XXIII, Bergamo, Italy
Vittoria	Fotia	ASST Papa Giovanni XXIII, Bergamo, Italy
Lorenzo	Chiudinelli	ASST Papa Giovanni XXIII, Bergamo, Italy
Michela	Franchi	ASST Papa Giovanni XXIII, Bergamo, Italy
Michela	Libertini	Fondazione Poliambulanza Istituto Ospedaliero, Brescia, Italy
Rossella	Bertulli	Fondazione IRCCS, Istituto Nazionale Tumori, Milano, Italy
Salvatore	Provenzano	Fondazione IRCCS, Istituto Nazionale Tumori, Milano, Italy
Daniele	Generali	ASST di Cremona, Cremona, Italy
Salvatore	Grisanti	Azienda Ospedaliera Spedali Civili, Brescia, Italy
Alice	Baggi	Azienda Ospedaliera Spedali Civili, Brescia, Italy
Valeria	Tovazzi	Azienda Ospedaliera Spedali Civili, Brescia, Italy
Corrado	Ficorella	Università degli Studi dell'Aquila, L'Aquila, Italy
Giampiero	Porzio	Università degli Studi dell'Aquila, L'Aquila, Italy
Alessandro	Parisi	Università degli Studi dell'Aquila, L'Aquila, Italy
Paola	Queirolo	Istituto Europeo di Oncologia, Milano, Italy
Maristella	Saponara	Istituto Europeo di Oncologia, Milano, Italy
Raffaele	Giusti	Azienda Ospedaliera Universitaria Sant'Andrea, Roma, Italy
Marco	Filetti	Azienda Ospedaliera Universitaria Sant'Andrea, Roma, Italy
Francesca	Mazzoni	Azienda Ospedaliera Universitaria Careggi, Firenze, Italy
Federica	Zoratto	Ospedale Santa Maria Goretti, Latina, Italy
Marco	Tucci	Policlinico Universitario di Bari, Bari, Italy
Rossana	Berardi	Università Politecnica delle Marche, Ancona, Italy
Luca	Cantini	Università Politecnica delle Marche, Ancona, Italy
Annalisa	Guida	Azienda Ospedaliera Santa Maria, Terni, Italy
Sergio	Bracarda	Azienda Ospedaliera Santa Maria, Terni, Italy
Clara	Martinez-Vila	Fundació Althaia Manresa, Manresa,
Maria	Iglesias	Hospital Son Llatzer Palma de Mallorca, Spain
Ana	Sanchez de Torre	Hospital XII de Octubre Madrid, Spain
Matteo	Lambertini	IRCCS Ospedale Policlinico San Martino, Genova, Italy; University of Genova, Genova, Italy.

Marco	Tagliamento	IRCCS Ospedale Policlinico San Martino, Genova, Italy; University of Genova, Genova, Italy.
-------	-------------	---

Figure 2 Legend:

Histogram reporting the corresponding prevalence of COVID-19 sequelae, hospital admission due to COVID-19 and complicated COVID-19 across the 6 defined age categories. **Sequelae:** <40 yo: 8/112 (7.1%), 40-49 yo: 12/158 (7.6%), 50-59 yo: 45/334 (13.5%), 60-69 yo: 82/370 (22.2%), 70-79 yo: 63/396 (15.9%), ≥ 80 yo: 24/180 (13.3%). **Hospital admission (patients already hospitalized excluded):** <40 yo: 33/88 (37.6%), 40-49 yo: 51/125 (40.8%), 50-59 yo: 135/278 (48.5%), 60-69 yo: 191/302 (63.2%), 70-79 yo: 205/306 (66.9%), ≥ 80 yo: 95/126 (75.4%). **Complicated:** <40 yo: 11/112 (9.8%), 40-49 yo: 27/158 (17.1%), 50-59 yo: 68/334 (20.4%), 60-69 yo: 110/370 (29.7%), 70-79 yo: 128/396 (32.3%), ≥ 80 yo: 58/180 (32.2%). Linear trends were tested with the Cochran–Armitage test. Yo; years old.